

Using NOGAPS Singular Vectors to Diagnose Large-scale Influences on Tropical Cyclogenesis

PI: Prof. Sharanya J. Majumdar

RSMAS/MPO, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149.

Phone: (305) 421 4779 Fax: (305) 421 4696 Email: smajumdar@rsmas.miami.edu

Co-PI: Dr Melinda S. Peng

Marine Meteorology Division, Naval Research Laboratory, 7 Grace Hopper Avenue, Monterey, CA 93943.

Phone: (831) 656 4704. Email: melinda.peng@nrlmry.navy.mil

Co-PI: Dr Carolyn A. Reynolds

Marine Meteorology Division, Naval Research Laboratory, 7 Grace Hopper Avenue, Monterey, CA 93943.

Phone: (831) 656 4728. Email: carolyn.reynolds@nrlmry.navy.mil

Co-PI: Prof. Chun-Chieh Wu

Dept of Atmospheric Sciences, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd, Taipei 10673, Taiwan.

Phone: +886-2-2363-2303. Email: cwu@typhoon.as.ntu.edu.tw

Grant Number: N000140810250

http://www.nrlmry.navy.mil/adap-bin/tcs_adap.cgi

LONG-TERM GOALS

The overarching goal is to provide an improved understanding of the synoptic-scale influences on tropical cyclone formation in the NW Pacific Ocean basin, in the context of error growth in forecast models. Benefits to the Navy would include improved forecast skill of developing tropical cyclones.

OBJECTIVES

To connect NOGAPS Singular Vector perturbation growth to synoptic-scale dynamical influences on tropical cyclone formation and structure change; and to evaluate the influence of assimilating TCS-08 observations into the Navy assimilation and forecast system, and investigate whether NOGAPS SV perturbations capture this influence.

APPROACH

Four significant tropical cyclones (TCs) formed during the TCS-08 field phase: Nuri, Sinlaku, Hagupit and Jangmi. The first stage of the research is the diagnosis of NOGAPS SVs associated with these storms. The real-time NOGAPS SV sensitivity guidance that was produced for developing systems in a region close to Guam (Fig. 1) during TCS-08 is currently being analyzed. If necessary, the NOGAPS SV guidance will be rerun by Co-PI Reynolds with a specific focus on verification regions centered on the developing systems. The dominant sensitivity patterns, evolving horizontal and vertical structure

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2008		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Using NOGAPS Singular Vectors To Diagnose Large-scale Influences On Tropical Cyclogenesis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Miami, RSMAS/MPO,,4600 Rickenbacker Causeway,Miami,FL,33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT The overarching goal is to provide an improved understanding of the synoptic-scale influences on tropical cyclone formation in the NW Pacific Ocean basin, in the context of error growth in forecast models. Benefits to the Navy would include improved forecast skill of developing tropical cyclones.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

and error growth will be investigated for these cases, from the formation stage through to typhoon status. The connection between these perturbation structures and synoptic-scale processes influencing developing tropical cyclones will be investigated, expanding upon the case study of Yamaguchi et al. (2008) for a single typhoon. In addition to the regular dry SVs, moist SVs will also be computed and analyzed for interesting cases.

The second stage of the project is to evaluate the effectiveness of dropwindsonde observations on NOGAPS predictions of tropical cyclone formation and structure change, via “data denial” in the NOGAPS/NAVDAS framework. The dropwindsondes from the USAF C-130, NRL P-3, DOTSTAR and Falcon aircraft will be used, as will those released from driftsonde balloons in areas of high SV sensitivity. The modification to the NOGAPS synoptic fields due to the special TCS-08 and T-PARC observations will be analyzed and compared with the SV structures, elucidating the effectiveness of the SV structures in accounting for perturbation growth in global models.




FIGURE 1. NOGAPS Singular Vector sensitivity guidance for system TCS033, which developed into Typhoon Sinlaku. The NOGAPS SVs were initiated on 00 UTC September 9th 2008, and SV growth between 00 UTC 10th and 00 UTC 12th was optimized. (a) integrated total energy of the leading 3 initial-time SVs; (b) sensitivity to vorticity in each of the first 3 SVs (denoted by number on map); (c) sensitivity to temperature in each of the first 3 SVs.

WORK COMPLETED

In Year 1, the primary task was to provide real-time NOGAPS SV sensitivity guidance for targeted observations during the TCS-08 field phase. Co-PI Reynolds provided this guidance daily (and sometimes twice daily) for a variety of verification regions, encompassing formation, structure change, recurvature and extratropical transition. Daily summaries of the SV guidance were prepared under the “Weather Targeting Blog” section on the EOL catalog. For example, Figure 1 illustrates the overall sensitivity for Sinlaku being associated with the subtropical ridge, with sensitivity to mid-lower level vorticity perturbations on the south-east side, and mid-level temperature perturbations towards the east of Sinlaku. Prof. Majumdar and his new graduate student, Munehiko Yamaguchi, spent 3 and 2 weeks respectively at the TCS-08 Operations Center in Monterey. When a storm was present, they coordinated the Elluminate sessions on targeted observations, synthesizing a wide array of products on predictability and targeting to facilitate decisions on aircraft deployment. The presentations accompanying these sessions are archived in the “Weather Targeting Blog” section on the EOL catalog. These products are now being reviewed retrospectively by the U.Miami and NRL Monterey teams to identify key cases for in-depth investigation in Years 2 and 3.

An undergraduate student at the University of Miami, Peter Finocchio was paid \$1000 from this grant to investigate multi-model ensemble forecasts, and their uncertainty in capturing TC formation and track. A paper will be submitted within the next year. Via matching salary for his summer work and a travel grant from U. Miami, Peter spent 17 days at the TCS-08 Operations Center to assist the TCS-08 team with weather briefings.

Additionally, Dr Eric Rappin, an assistant scientist funded partially by a different ONR TCS-08 grant (N000140810251; PI Velden) has been training Majumdar and Yamaguchi on how to run a 2-km resolution version of the Weather Research and Forecasting (WRF) model, including a bogus vortex initialization. This high-resolution model can be nested within NOGAPS output to answer questions about tropical cyclone *structure change*.

RESULTS

Some general conclusions can be drawn via inspections of the NOGAPS SV guidance for mature tropical cyclones. For example, sensitivity is usually on the eastern and southern side of the cyclone, corresponding to the periphery of the subtropical ridge influencing the cyclone's motion and wind field, and to the north and north-west, associated with the interaction with an approaching shortwave trough. These results are consistent with those found in earlier studies by Co-PIs Peng and Reynolds. However, for formation, which is the primary and most novel focus of this project, the cases need to be investigated in more detail since the sensitivity appears less intuitive on first inspection.

IMPACT/APPLICATIONS

The scientific impact will be an improved understanding of the underlying environmental mechanisms that influence tropical cyclone formation. This understanding will be coupled with a quantitative knowledge of error growth in global models, via SVs. The SVs also possess practical value in that they can be used in future targeting applications, with the augmentation of the routine observational network with new aircraft, balloon and satellite-based observations to improve Navy forecasts of typhoon formation and structure change.

RELATED PROJECTS

This project is related to that funded by the TCS-08 grant N000140810251: "Advanced Satellite-Derived Wind Observations, Assimilation, and Targeting Strategies during TCS-08 for Developing Improved Operational Analysis and Prediction of Western North Pacific Tropical Cyclones", on which Majumdar is a Co-PI. The NOGAPS Singular Vectors will also be used as a targeting strategy in this TCS-08 grant.

PUBLICATIONS

Yamaguchi, M., T. Iriguchi, T. Nakazawa and C.-C. Wu, 2008: An observing system experiment for Typhoon Conson (2004) using a singular vector method and DOTSTAR data, *Mon. Wea. Rev.*, accepted.